



Dieldrin in cod liver from the North Sea - Time trend from 1998

Fromberg, Arvid; Højgård, Arne; Poulsen, Mette Erecius

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Fromberg, A., Højgård, A., & Poulsen, M. E. (2012). *Dieldrin in cod liver from the North Sea - Time trend from 1998*. Poster session presented at 9th European Pesticide Residue Workshop , Vienna, Austria.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Dieldrin in cod liver from the North Sea - Time trend from 1988

Arvid Fromberg¹, Arne Højgård² and Mette Erecius Poulsen¹

1. DTU National Food Institute, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark; e-mail: arfr@food.dtu.dk
2. Regional Veterinary and Food Administration Centre, Region West, Sønderskovvej 5, DK-8520 Lystrup, Denmark,

Monitoring of pesticide levels in fish samples is an important contribution to the intake estimations



Background for the monitoring programme

Monitoring programs for pesticides in food often include investigations of persistent organochlorine contaminants and metabolites and are carried out because of their potentially hazardous health effects on humans. Due to the organochlorine pesticides stability and tendency to accumulate in fatty tissues they can enter the food chain and therefore be found in marine species. Relatively high levels occur in cod liver, which has high lipid content and for that reason is suitable for monitoring purposes. Results for dieldrin in cod caught in the North Sea and Skagerrak from 1988 to 2010 are presented looking at the time trend and decline in the levels during the period including 116 samples.

Estimated intake of dieldrin
from consumption of fish

Mean	0.035
Median	0.015
90% fractile	0.093
95% fractile	0.125

Results give as µg/day




Adults, estimated intake from fish

The location of the North Sea

Levels of dieldrin in cod liver has decreased during the last decades from 1988, however is this still the case for the North Sea and Skagerrak?

Time trend for dieldrin

Contents of organochlorine environmental contaminants in fish have decreased since 1988; however developments in recent years do not show a clear trend, but rather a more or less steady state condition. Statistical analyses of data from cod liver showed that the distribution of concentrations is best described by a logarithmic normal distribution, and that the development over time can be described by a linear regression based on logarithmized data.



Time trend for the level of dieldrin in cod liver from fish caught in the North Sea and Skagerrak